Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to 508 standards due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehp508@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Comparing the Health Effects of Ambient Particulate Matter Estimated Using Ground-Based versus Remote Sensing Exposure Estimates

Michael Jerrett, Michelle C. Turner, Bernardo S. Beckerman, C. Arden Pope III, Aaron van Donkelaar, Randall V. Martin, Marc Serre, Dan Crouse, Susan M. Gapstur, Daniel Krewski, W. Ryan Diver, Patricia F. Coogan, George D. Thurston, and Richard T. Burnett

Table of Contents

Methods. Detailed Description of the Individual and Ecological Variables included in the Cox proportional Hazards Models

- **Table S1.** Distribution (n, %) of selected participant characteristics at enrollment (1982), ACS CPS II cohort, U.S., compared to those excluded from the analysis with commentary on the differences
- **Table S2.** Results of the Cox Proportional Hazard Modeling with adjustment for individual or individual plus year 1990 ecologic covariates. Hazard ratios expressed over an IDR increment.
- **Table S3.** Descriptive Statistics for the Central Monitor Exposure Estimates.
- **Table S4.** Results of the Cox Proportional Hazard Modeling with adjustment for individual plus year 1990 ecologic covariates plus MSA size or elevation or RE model. Hazard ratios expressed over a $10 \, \mu g/m3$ increment for diseases of the circulatory system.
- **Table S5.** Results of the Cox Proportional Hazard Modeling with adjustment for

individual plus year 1990 ecologic covariates plus MSA size or elevation or RE model. Hazard ratios expressed over a 10 µg/m3 increment for ischemic heart disease.

Table S6. Results of the Cox Proportional Hazard Modeling with adjustment for individual or individual plus year 2000 ecologic covariates. Hazard ratios expressed over a 10 μg/m3 increment.

Table S7. Results of the Cox proportional hazard modeling with adjustment for individual or individual plus year 1990 ecologic covariates for mortality from diabetes. Hazard ratios expressed over a 10 μg/m³ increment.

Figure S1. Fully-Adjusted hazard ratios (HR) from Table 2 for deaths of the circulatory system and IHD with HR plotted against AIC statistics to illustrate the effect of model fit on the HR among seven exposure models.

Figure S2. Concentration response plot for PM2.5 BMELUR 02-04 with two degrees of freedom for circulatory mortality

Figure S3. Concentration response plot for PM2.5 RS no GWR CT 02-04 with two degrees of freedom for circulatory mortality